


Fractional dose IPV

Polio type 2 withdrawal


Polio Eradication : Endgame plans

Epidemiology Unit




- Detect and interrupt all poliovirus transmission
- Strengthen immunization systems and withdraw oral polio vaccine
- Contain poliovirus and certify interruption of transmission
- Plan polio's legacy

Polio Endgame Objective 2 : three distinct stages



Plan of OPV withdrawal in Polio Eradication



fIPV

- 0.1 ml intradermal injection (10-15° angle)
- Left arm, Deltoid region
- 2 doses
- 2nd and 4th months of the age (together with OPV and Pentavalent)
 - Pentavalent to left thigh
 - fIPV (0.1 ml) intradermal to left arm



Step 1: OPV



Step 2: Penta
(left thigh)



Step 3: fIPV
(left arm)

Data recording

- CHDR
 - Other vaccine section
 - 2 months : fIPV1
 - 4 months: fIPV2
- Clinic Immunization Register (H-1216)
 - Divide IPV column in to 2 and mention (fIPV 1, fIPV 2)
- Clinic Summary (RH-MIS 518)
 - Divide IPV column in to 2 and mention (fIPV 1, fIPV 2)
- Birth & Immunization Register(EPI/03/79-2014)
 - Divide IPV column in to 2 and mention (fIPV 1, fIPV 2)
- WEBIIS: necessary changes will be done for data entry

Intradermal vaccination procedure

