



# WEEKLY EPIDEMIOLOGICAL REPORT

A publication of the Epidemiology Unit  
Ministry of Healthcare and Nutrition

231, de Saram Place, Colombo 01000, Sri Lanka  
Tele: + 94 11 2695112, Fax: +94 11 2696583, E mail: epidunit@sltnet.lk  
Epidemiologist: +94 11 2681548, E mail: chepid@sltnet.lk  
Web: <http://www.epid.gov.lk>

Vol. 37 No.49

04<sup>th</sup> – 10<sup>th</sup> December 2010

## Crimean - Congo Haemorrhagic fever

Crimean-Congo haemorrhagic fever (CCHF) is a viral haemorrhagic fever of the *Nairovirus* group. Although primarily a zoonosis, sporadic cases and outbreaks of CCHF affecting humans do occur. The disease is endemic in many countries in Africa, Europe and Asia, and during 2001, cases or outbreaks have been recorded in Kosovo, Albania, Iran, Pakistan, and South Africa.

The disease was first described in the Crimea in 1944 and given the name Crimean haemorrhagic fever. In 1969 it was recognized that the pathogen causing Crimean haemorrhagic fever was the same as that responsible for an illness identified in 1956 in the Congo, and linkage of the 2 place names resulted in the current name for the disease and the virus. CCHF is a severe disease in humans, with a high mortality rate. Fortunately, human illness occurs infrequently, although animal infection may be more common.

The geographical distribution of the virus, like that of its tick vector, is widespread. Evidence of CCHF virus has been found in Africa, Asia, the Middle East and Eastern Europe. Healthcare workers in endemic areas should be aware of the illness and the correct infection control procedures to protect themselves and their patients from the risk of nosocomial (hospital-acquired) infection.

### CCHF virus

The virus which causes CCHF is a *Nairovirus*, a group of related viruses forming one of the five genera in the *Bunyaviridae* family of viruses. All of the 32 members of the *Nairovirus* genus are transmitted by argasid or ixodid ticks, but only three have been implicated as causes of human disease: the Dugbe and Nairobi sheep viruses, and CCHF, which is the

most important human pathogen amongst them.

### CCHF reservoirs and vectors

The CCHF virus may infect a wide range of domestic and wild animals. Many birds are resistant to infection, but ostriches are susceptible and may show a high prevalence of infection in endemic areas. Animals become infected with CCHF from the bite of infected ticks.

A number of tick genera are capable of becoming infected with CCHF virus, but the most efficient and common vectors for CCHF appear to be members of the *Hyalomma* genus. Trans-ovarial (transmission of the virus from infected female ticks to offspring via eggs) and venereal transmission have been demonstrated amongst some vector species, indicating one mechanism which may contribute to maintaining the circulation of the virus in nature.

However, the most important source for acquisition of the virus by ticks is believed to be infected small vertebrates on which immature *Hyalomma* ticks feed. Once infected, the tick remains infected through its developmental stages, and the mature tick may transmit the infection to large vertebrates, such as livestock. Domestic ruminant animals, such as cattle, sheep and goats, are viraemic (virus circulating in the bloodstream) for around one week after becoming infected.

Humans who become infected with CCHF acquire the virus from direct contact with blood or other infected tissues from livestock during this time, or they may become infected from a tick bite. The majority of cases have occurred in those involved with the livestock industry, such as agricultural workers, slaughterhouse workers and veterinarians.

Contents	Page
1. Article : Crimean Congo Haemorrhagic Fever	1
2. Surveillance of vaccine preventable diseases & AFP (27 <sup>th</sup> November– 03 <sup>rd</sup> December 2010)	3
3. Summary of newly introduced notifiable diseases (27 <sup>th</sup> November– 03 <sup>rd</sup> December 2010)	3
4. Summary of selected notifiable diseases reported (27 <sup>th</sup> November– 03 <sup>rd</sup> December 2010)	4

### Clinical features

The length of the incubation period for the illness appears to depend on the mode of acquisition of the virus. Following infection via tick bite, the incubation period is usually one to three days, with a maximum of nine days. The incubation period following contact with infected blood or tissues is usually five to six days, with a documented maximum of 13 days.

Onset of symptoms is sudden, with fever, myalgia (aching muscles), dizziness, neck pain and stiffness, backache, headache, sore eyes and photophobia (sensitivity to light). There may be nausea, vomiting and sore throat early on, which may be accompanied by diarrhoea and generalised abdominal pain. Over the next few days, the patient may experience sharp mood swings, and may become confused and aggressive. After two to four days, the agitation may be replaced by sleepiness, depression and lassitude, and the abdominal pain may localize to the right upper quadrant, with detectable hepatomegaly (liver enlargement).

Other clinical signs which emerge include tachycardia (fast heart rate), lymphadenopathy (enlarged lymph nodes), and a petechial rash (a rash caused by bleeding into the skin), both on internal mucosal surfaces, such as in the mouth and throat, and on the skin. The petechiae may give way to ecchymoses (like a petechial rash, but covering larger areas) and other haemorrhagic phenomena such as melaena (bleeding from the upper bowel, passed as altered blood in the faeces), haematuria (blood in the urine), epistaxis (nosebleeds) and bleeding from the gums. There is usually evidence of hepatitis. The severely ill may develop hepatorenal (i.e., liver and kidney) and pulmonary failure after the fifth day of illness.

The mortality rate from CCHF is approximately 30%, with death occurring in the second week of illness. In those patients who recover, improvement generally begins on the ninth or tenth day after the onset of illness.

### Diagnosis

Diagnosis of suspected CCHF is performed in specially-equipped, high biosafety level laboratories. IgG and IgM antibodies may be detected in serum by enzyme-linked immunoassay (the "ELISA" or "EIA" methods) from about day six of illness. IgM remains detectable for up to four months, and IgG levels decline but remain detectable for up to five years.

Patients with fatal disease do not usually develop a measurable antibody response and in these individuals, as well as in patients in the first few days of illness, diagnosis is achieved by virus detection in blood or tissue samples. There are several methods for doing this. The virus may be isolated from blood or tissue specimens in the first five days of illness, and grown in cell culture. Viral antigens may sometimes be shown in tissue samples using immunofluorescence or EIA.

More recently, the polymerase chain reaction (PCR), a molecular method for detecting the viral genome, has been successfully applied in diagnosis.

### Treatment

- General supportive therapy is the mainstay of patient management in CCHF. Intensive monitoring to guide volume and blood component replacement is required.
- The antiviral drug ribavirin has been used in treatment of established CCHF infection with apparent benefit. Both oral and intravenous formulations seem to be effective.
- The value of immune plasma from recovered patients for therapeutic purposes has not been demonstrated, although it has been employed on several occasions.

### Prevention and control

- Although an inactivated, mouse brain-derived vaccine against CCHF has been developed and used on a small scale in Eastern Europe, there is no safe and effective vaccine widely available for human use. The tick vectors are numerous and widespread and tick control with acaricides (chemicals intended to kill ticks) is only a realistic option for well-managed livestock production facilities.
- Persons living in endemic areas should use personal protective measures that include avoidance of areas where tick vectors are abundant and when they are active; regular examination of clothing and skin for ticks, and their removal; and use of repellents.
- Persons who work with livestock or other animals in the endemic areas can take practical measures to protect themselves. These include the use of repellents on the skin (e.g. DEET) and clothing (e.g. permethrin) and wearing gloves or other protective clothing to prevent skin contact with infected tissue or blood.
- When patients with CCHF are admitted to hospital, there is a risk of nosocomial spread of infection. In the past, serious outbreaks have occurred in this way and it is imperative that adequate infection control measures be observed to prevent this disastrous outcome.
- Patients with suspected or confirmed CCHF should be isolated and cared for using barrier nursing techniques. Specimens of blood or tissues taken for diagnostic purposes should be collected and handled using universal precautions. Sharps (needles and other penetrating surgical instruments) and body wastes should be safely disposed of using appropriate decontamination procedures.
- Healthcare workers are at risk of acquiring infection from sharps injuries during surgical procedures and, in the past, infection has been transmitted to surgeons operating on patients to determine the cause of the abdominal symptoms in the early stages of (at that moment undiagnosed) infection. Healthcare workers who have had contact with tissue or blood from patients with suspected or confirmed CCHF should be followed up with daily temperature and symptom monitoring for at least 14 days after the putative exposure.

Source: WHO

**Table 1: Vaccine-preventable Diseases & AFP**

27<sup>th</sup> November – 03<sup>rd</sup> December 2010(48<sup>th</sup> Week)

Disease	No. of Cases by Province									Number of cases during current week in 2010	Number of cases during same week in 2009	Total number of cases to date in 2010	Total number of cases to date in 2009	Difference between the number of cases to date in 2010 & 2009
	W	C	S	N	E	NW	NC	U	Sab					
Acute Flaccid Paralysis	00	00	00	00	00	00	00	00	00	00	02	77	68	+ 13.2 %
Diphtheria	00	00	00	00	00	00	00	00	00	00	00	00	00	-
Measles	00	00	00	00	00	00	00	00	00	00	03	88	165	- 46.7 %
Tetanus	00	00	00	00	00	00	00	00	00	00	00	22	27	- 18.5 %
Whooping Cough	00	00	00	00	00	00	00	00	00	00	00	30	61	- 50.8 %
Tuberculosis	24	03	22	03	15	23	00	10	24	124	436	9435	9771	-03.4 %

**Table 2: Newly Introduced Notifiable Disease**

27<sup>th</sup> November – 03<sup>rd</sup> December 2010(48<sup>th</sup> Week)

Disease	No. of Cases by Province									Number of cases during current week in 2010	Number of cases during same week in 2009	Total number of cases to date in 2010	Total number of cases to date in 2009	Difference between the number of cases to date in 2010 & 2009
	W	C	S	N	E	NW	NC	U	Sab					
Chickenpox	13	06	06	00	03	08	00	04	06	46	57	3176	14036	- 77.4 %
Meningitis	09 CB=3 KL=3 GM=3	01 MT=1	01 GL=1	00	04 TR=1 BT=3	00	05 AP=4 PO=1	01 MO=1	01 KG=1	22	32	1483	1546	- 4.1 %
Mumps	5	6	4	1	1	11	9	1	9	47	21	1149	1640	- 29.9 %
Leishmaniasis	00	00	12 HB=12	00	00	00	01 AP=1	00	00	13	04	381	636	- 40.1 %

**Key to Table 1 & 2**

**Provinces:** W: Western, C: Central, S: Southern, N: North, E: East, NC: North Central, NW: North Western, U: Uva, Sab: Sabaragamuwa.

**DPDHS Divisions:** CB: Colombo, GM: Gampaha, KL: Kalutara, KD: Kandy, ML: Matale, NE: Nuwara Eliya, GL: Galle, HB: Hambantota, MT: Matara, JF: Jaffna, KN: Killinochchi, MN: Mannar, VA: Vavuniya, MU: Mullaitivu, BT: Batticaloa, AM: Ampara, TR: Trincomalee, KM: Kalmunai, KR: Kurunegala, PU: Puttalam, AP: Anuradhapura, PO: Polonnaruwa, BD: Badulla, MO: Moneragala, RP: Ratnapura, KG: Kegalle.

**Data Sources:**

**Weekly Return of Communicable Diseases:** Diphtheria, Measles, Tetanus, Whooping Cough, Chickenpox, Meningitis, Mumps.

**Special Surveillance:** Acute Flaccid Paralysis.

Leishmaniasis is notifiable only after the General Circular No: 02/102/2008 issued on 23 September 2008.

**Dengue Prevention and Control Health Messages**

**You have a duty and a responsibility in preventing dengue fever. Make sure that your environment is free from water collections where the dengue mosquito could breed**

**Table 4: Selected notifiable diseases reported by Medical Officers of Health**  
27<sup>th</sup> November – 03<sup>rd</sup> December 2010(48<sup>th</sup> Week)

DPDHS Division	Dengue Fever / DHF*		Dysentery		Encephalitis		Enteric Fever		Food Poisoning		Leptospirosis		Typhus Fever		Viral Hepatitis		Human Rabies		Returns received timely
	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	%
Colombo	14	5804	2	298	0	16	5	190	0	49	5	563	0	9	0	68	0	1	92
Gampaha	16	3886	4	181	1	28	2	65	0	21	17	563	0	16	0	125	0	6	73
Kalutara	4	1796	2	246	0	15	0	41	0	76	11	416	0	5	0	41	0	3	75
Kandy	5	1633	3	330	0	6	1	34	0	16	4	172	0	139	2	151	0	1	87
Matale	13	618	10	310	0	8	0	36	0	81	6	112	0	7	0	54	0	1	100
Nuwara Eliya	0	223	3	344	0	1	2	119	0	89	0	33	1	67	0	52	0	0	92
Galle	1	1106	1	250	1	9	0	14	0	59	10	165	2	25	0	22	0	5	68
Hambantota	0	799	5	93	0	7	0	4	1	16	2	100	1	92	1	20	0	0	82
Matara	5	610	2	168	1	9	0	14	0	53	6	372	1	136	0	19	0	0	88
Jaffna	4	2902	2	296	0	7	5	604	0	10	0	1	2	141	7	86	0	2	75
Kilinochc	1	48	0	18	0	0	0	10	0	1	0	3	0	0	0	1	0	1	100
Mannar	1	564	0	48	0	2	1	47	0	10	0	0	0	1	0	17	0	1	33
OVavuniya	0	575	1	59	0	3	0	45	0	13	0	2	0	1	0	13	0	2	75
Mullaitivu	0	20	0	8	0	0	0	3	0	0	0	0	0	0	0	1	1	1	50
Batticaloa	12	1232	2	201	0	4	1	40	0	38	0	13	0	4	0	7	0	4	86
Ampara	0	156	0	116	0	1	0	9	0	65	0	31	0	1	0	14	0	0	57
Trincomalee	9	974	4	160	0	14	0	7	0	14	3	41	0	21	1	16	0	1	82
Kurunegala	6	1406	18	366	0	20	4	59	0	35	9	368	0	59	0	127	0	4	95
Puttalam	4	1003	5	225	0	7	0	56	0	125	0	77	3	8	0	23	0	1	56
Anuradhapur	9	1077	4	155	0	11	0	16	0	46	0	96	0	30	0	53	0	4	74
Polonnaruwa	0	395	2	112	0	2	0	8	0	10	4	88	0	2	0	47	0	0	100
Badulla	6	1301	1	215	0	1	2	97	0	1029	0	88	1	119	0	111	0	0	80
Monaragala	1	1048	0	184	1	2	1	45	0	7	3	42	0	89	2	99	0	3	55
Ratnapura	5	2772	1	474	0	8	0	24	0	27	3	407	0	62	0	102	0	3	50
Kegalle	1	898	3	160	0	18	1	79	0	29	20	394	0	32	3	133	0	0	82
Kalmunai	0	545	3	312	0	3	1	14	0	9	0	3	0	0	0	13	0	1	62
<b>SRI LANKA</b>	<b>117</b>	<b>33391</b>	<b>78</b>	<b>5329</b>	<b>4</b>	<b>202</b>	<b>26</b>	<b>1680</b>	<b>01</b>	<b>928</b>	<b>103</b>	<b>4150</b>	<b>11</b>	<b>1066</b>	<b>16</b>	<b>1415</b>	<b>01</b>	<b>45</b>	<b>77</b>

Source: Weekly Returns of Communicable Diseases WRCD).

\*Dengue Fever / DHF refers to Dengue Fever / Dengue Haemorrhagic Fever.

\*\*Timely refers to returns received on or before 03<sup>rd</sup> December, 2010 Total number of reporting units =320. Number of reporting units data provided for the current week: 249

A = Cases reported during the current week. B = Cumulative cases for the year.

**PRINTING OF THIS PUBLICATION IS FUNDED BY THE WORLD HEALTH ORGANIZATION (WHO).**

Comments and contributions for publication in the WER Sri Lanka are welcome. However, the editor reserves the right to accept or reject items for publication. All correspondence should be mailed to The Editor, WER Sri Lanka, Epidemiological Unit, P.O. Box 1567, Colombo or sent by E-mail to [chepid@slt.net.lk](mailto:chepid@slt.net.lk).

**ON STATE SERVICE**

**Dr. P. PALIHAWADANA**  
CHIEF EPIDEMIOLOGIST  
EPIDEMIOLOGY UNIT  
231, DE SARAM PLACE  
COLOMBO 10