



WEEKLY EPIDEMIOLOGICAL REPORT

A publication of the Epidemiology Unit
Ministry of Health

231, de Saram Place, Colombo 01000, Sri Lanka
Tele: + 94 11 2695112, Fax: +94 11 2696583, E mail: epidunit@slt.net.lk
Epidemiologist: +94 11 2681548, E mail: chepid@slt.net.lk
Web: <http://www.epid.gov.lk>

Vol. 42 No. 37

05th – 11th September 2015

Toxocariasis

Toxocariasis is an infection transmitted from animals to humans (zoonosis) caused by the parasitic roundworms commonly found in the intestine of dogs (*Toxocara canis*) and cats (*T. cati*).

Epidemiology & Risk Factors

Infected dogs and cats shed *Toxocara* eggs in their faeces into the environment. Once in the environment, it takes 2 to 4 weeks for *Toxocara* larvae to develop and for the eggs to become infectious. Humans or other animals can be infected by accidentally ingesting *Toxocara* eggs. For example, humans can become infected if they work with dirt and accidentally ingest dirt containing *Toxocara* eggs. Although rare, people can be infected by eating undercooked or raw meat from an infected animal such as a lamb or rabbit. Because dogs and cats are frequently found where people live, there may be large numbers of infected eggs in the environment. Once in the body, the *Toxocara* eggs hatch and roundworm larvae can travel in the bloodstream to different parts of the body, including the liver, heart, lungs, brain, muscles, or eyes. Most infected people do not have any symptoms. However, in some people, the *Toxocara* larvae can cause damage to these tissues and organs. The symptoms of toxocariasis, the disease

caused by these migrating larvae, include fever, coughing, inflammation of the liver or eye problems.

A U.S. study in 1996 showed that 30% of dogs younger than 6 months deposit *Toxocara* eggs in their feces; other studies have shown that almost all puppies are born already infected with *Toxocara canis*. Research also suggests that 25% of all cats are infected with *Toxocara cati*. Infection rates are higher for dogs and cats that are left outside for more time and allowed to eat other animals. Globally, toxocariasis is found in many countries, and prevalence rates can reach as high as 40% or more in parts of the world. There are several factors that have been associated with higher rates of infection with *Toxocara*. People are more likely to be infected with *Toxocara* if they own a dog. In Sri Lanka, sero-prevalence studies showed that *Toxocara* sero-positivity in children with asthma was 29% and the proportion of children who were positive for *Toxocara* antibodies in the study population was 20%. Children and adolescents under the age of 20 are more likely to test positive for *Toxocara* infection. This may be because children are more likely to eat dirt and play in outdoor environments, such as sandboxes, where dog and cat faeces can be found. This

Contents

Page

- | | |
|---|---|
| 1. <i>Leading Article – Toxocariasis</i> | 1 |
| 2. <i>Summary of selected notifiable diseases reported - (29th – 04th September 2015)</i> | 3 |
| 3. <i>Surveillance of vaccine preventable diseases & AFP - (29th – 04th September 2015)</i> | 4 |

infection is more common in people living in poverty. Geographic location plays a role as well, because Toxocara is more prevalent in hot, humid regions where eggs are kept viable in the soil.

Disease

Most people infected with Toxocara do not have any symptoms. There are two major forms of toxocariasis, visceral toxocariasis (VT), also called visceral larva migrans (VLM), and ocular toxocariasis (OT), also called ocular larva migrans (OLM). The syndromes VLM and OLM can be caused by infection with the migrating larvae of other kinds of parasites which cause symptoms similar to those caused by migrating Toxocara larvae.

In a few people who are infected with high numbers of Toxocara larvae or have repeated infections, the larvae can travel through parts of the body such as the liver, lungs, or central nervous system and cause symptoms such as fever, coughing, enlarged liver or pneumonia. This form of toxocariasis is called visceral toxocariasis (VT). The larvae can also travel to the eye and cause ocular toxocariasis (OT). Ocular toxocariasis occurs when a microscopic Toxocara larva enters the eye and causes inflammation and scarring on the retina. OT typically occurs only in one eye and can cause irreversible vision loss.

Diagnosis

Diagnosis of either visceral toxocariasis or ocular toxocariasis are based on the presence of signs of VT or OT and history of exposure to a potential source of infectious Toxocara eggs. The diagnosis of visceral toxocariasis is based on compatible disease and exposure history with positive results by serological testing. Currently recommended test is an enzyme-linked immunosorbent assay (ELISA) with larval stage antigens. Usually excretory/secretory antigens are released when infective Toxocara larvae are cultured. The specificity of this assay is good although cross reactivity with antibody to the human roundworm, *Ascaris lumbricoides*, is possible; however, assays employing Toxocara excretory/secretory antigens minimize this problem. Positive serological results should be interpreted with consideration of the patient's clinical

status. Detectable antibody may be the result of infection in the past. Also, seropositivity can be present in asymptomatic Toxocara infection. Paired serum samples demonstrating a significant rise in antibody level over time may be useful to confirm active infection.

Treatment

Albendazole-400 mg by mouth twice a day for five days (both adult and pediatric dosage)

Mebendazole-100-200 mg by mouth twice a day for five days (both adult and pediatric dosage)

Prevention & Control

Controlling Toxocara infection in dogs and cats will reduce the number of infectious eggs in the environment and reduce the risk of infection for people.

- Clean the pet's living area at least once a week. Faeces should be either buried or bagged and disposed of in the trash. Wash hands after handling pet waste.
- Do not allow children to play in areas that are soiled with pet or other animal faeces and cover sandboxes when not in use to make sure that animals do not get inside and contaminate them.
- Wash hands with soap and warm water after playing with your pets or other animals, after outdoor activities and before handling food.
- Teach children the importance of washing hands to prevent infection.

Sources

1. Toxocara seropositivity in Sri Lankan children with asthma available at http://www.researchgate.net/publication/24392974-Toxocare_-_Seropositivity_in_Sri-Lankan_Children_With_asthma
2. Toxocariasis, available at <http://www.cdc.gov/parasites/toxocariasis/>

Compiled by Dr.H.H.W.S.B Herath of the Epidemiology Unit

Table 1: Selected notifiable diseases reported by Medical Officers of Health 29th - 04th Sep 2015 (36th Week)

RDHS Division	Dengue Fever		Dysentery		Encephalitis		Enteric Fever		Food Poisoning		Leptospirosis		Typhus Fever		Viral Hepatitis		Human Rabies		Chickenpox		Meningitis		Leishmaniasis		WRCD	
	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B	T*	C**
Colombo	116	6436	0	139	1	9	4	74	3	104	3	207	1	9	1	28	0	3	3	344	3	30	0	0	94	6
Gampaha	48	2816	1	66	0	6	1	26	0	25	7	272	1	9	4	108	0	0	11	192	0	19	0	2	93	7
Kalutara	24	1035	2	74	0	4	0	33	0	72	3	224	0	3	2	27	0	2	3	219	1	37	0	0	100	0
Kandy	11	828	3	88	0	6	0	27	4	38	1	85	1	50	0	107	0	0	8	171	2	14	0	11	100	0
Matale	0	341	0	35	0	1	0	8	0	5	1	50	0	8	1	26	0	0	1	21	1	19	0	15	54	46
NuwaraEliya	2	118	6	262	0	3	3	18	0	7	1	29	1	49	0	43	0	0	1	100	2	43	0	1	92	8
Galle	11	551	0	60	0	3	0	7	0	19	1	168	2	57	0	7	0	0	4	201	2	40	0	2	80	20
Hambantota	9	228	0	24	0	1	0	8	0	24	2	71	1	40	0	27	0	0	2	90	0	11	0	219	92	8
Matara	4	282	0	51	0	6	0	4	0	44	9	125	0	27	2	28	0	0	3	185	0	16	5	95	100	0
Jaffna	14	1250	20	622	0	9	1	158	2	65	0	14	0	540	0	10	0	2	0	164	0	15	0	0	92	8
Kilinochchi	0	54	0	67	0	0	0	13	0	31	0	1	0	21	0	0	0	1	0	15	0	0	0	0	50	50
Mannar	0	77	0	9	0	1	0	5	0	3	0	8	0	20	0	0	0	0	0	7	0	0	0	1	80	20
Vavuniya	1	93	0	16	0	6	0	57	0	11	0	17	0	13	0	1	0	2	1	37	0	12	0	6	75	25
Mullaitivu	1	114	0	23	0	2	1	12	0	1	0	5	0	9	0	3	0	1	0	5	0	3	0	5	40	60
Batticaloa	1	1320	7	248	1	7	0	25	0	139	0	12	0	3	0	11	0	1	1	45	0	16	0	0	64	36
Ampara	0	43	2	36	0	1	0	1	0	10	0	11	0	2	0	4	0	0	1	168	0	5	0	3	57	43
Trincomalee	2	513	1	45	0	0	0	30	0	35	0	14	1	20	1	8	0	1	1	74	0	6	0	2	75	25
Kurunegala	7	970	4	132	0	2	0	6	0	13	0	200	0	28	1	34	0	6	3	318	0	29	4	98	70	30
Puttalam	1	545	1	38	0	4	0	7	0	9	0	25	0	18	0	1	0	0	1	40	0	24	0	2	62	38
Anuradhapura	3	306	3	65	0	2	0	3	0	58	0	182	0	19	0	12	0	1	3	141	2	27	9	249	79	21
Polonnaruwa	1	156	0	29	0	4	0	9	0	3	0	64	0	1	0	4	0	0	1	97	1	20	0	73	43	57
Badulla	1	420	7	159	1	7	0	9	0	12	3	56	5	92	3	153	0	2	4	159	2	67	0	6	88	12
Monaragala	0	152	1	88	0	4	1	15	1	5	0	135	1	63	8	98	0	1	1	78	1	19	2	27	100	0
Ratnapura	10	758	1	226	2	14	0	37	0	8	3	246	1	52	1	173	0	1	3	101	0	45	0	15	67	33
Kegalle	4	432	2	55	1	9	2	59	0	9	3	229	0	38	0	71	0	0	1	165	0	40	0	0	64	36
Kalmunei	0	440	1	94	0	1	0	1	2	45	0	7	0	0	0	1	0	0	1	92	0	9	0	0	46	54
SRILANKA	271	20278	62	2751	6	112	13	652	12	795	37	2457	15	1191	24	985	0	24	58	3229	17	566	20	832	79	21

Source: Weekly Returns of Communicable Diseases (WRCD).

*T=Timeliness refers to returns received on or before 04th September, 2015 Total number of reporting units 337 Number of reporting units data provided for the current week: 268 C**=Completeness
A = Cases reported during the current week. B = Cumulative cases for the year.

Table 2: Vaccine-Preventable Diseases & AFP

29th - 04th Sep 2015 (36th Week)

Disease	No. of Cases by Province									Number of cases during current week in 2015	Number of cases during same week in 2014	Total number of cases to date in 2015	Total number of cases to date in 2014	Difference between the number of cases to date in 2014 & 2015
	W	C	S	N	E	NW	NC	U	Sab					
AFP*	00	00	00	00	00	00	01	00	01	02	01	53	60	-11.6%
Diphtheria	00	00	00	00	00	00	00	00	00	00	00	00	00	0%
Mumps	01	00	01	00	00	01	00	03	00	06	12	270	514	-47.4%
Measles	19	05	08	00	05	02	05	02	07	53	20	2093	2564	-18.3%
Rubella	00	00	00	00	00	00	00	00	00	00	01	08	15	-46.6%
CRS**	00	00	00	00	00	00	00	00	00	00	00	00	04	-100%
Tetanus	00	00	00	00	00	00	00	00	00	00	00	14	10	+40%
Neonatal Tetanus	00	00	00	00	00	00	00	00	00	00	00	00	00	0%
Japanese Encephalitis	00	00	00	00	00	00	00	00	00	00	01	07	22	-68.1%
Whooping Cough	01	00	00	00	00	00	00	00	00	01	01	63	40	+57.5%
Tuberculosis	60	18	18	17	17	00	00	02	12	144	132	6907	6768	+2.1%

Key to Table 1 & 2

Provinces: W: Western, C: Central, S: Southern, N: North, E: East, NC: North Central, NW: North Western, U: Uva, Sab: Sabaragamuwa.
 RDHS Divisions: CB: Colombo, GM: Gampaha, KL: Kalutara, KD: Kandy, ML: Matale, NE: Nuwara Eliya, GL: Galle, HB: Hambantota, MT: Matara, JF: Jaffna, KN: Killinochchi, MN: Mannar, VA: Vavuniya, MU: Mullaitivu, BT: Batticaloa, AM: Ampara, TR: Trincomalee, KM: Kalmunai, KR: Kurunegala, PU: Puttalam, AP: Anuradhapura, PO: Polonnaruwa, BD: Badulla, MO: Moneragala, RP: Ratnapura, KG: Kegalle.

Data Sources:

Weekly Return of Communicable Diseases: Diphtheria, Measles, Tetanus, Neonatal Tetanus, Whooping Cough, Chickenpox, Meningitis, Mumps., Rubella, CRS,

Special Surveillance: AFP* (Acute Flaccid Paralysis), Japanese Encephalitis

CRS** =Congenital Rubella Syndrome

AFP and all clinically confirmed Vaccine Preventable Diseases except Tuberculosis and Mumps should be investigated by the MOH

Dengue Prevention and Control Health Messages

Look for plants such as bamboo, bohemia, rampe and banana in your surroundings and maintain them

PRINTING OF THIS PUBLICATION IS FUNDED BY THE WORLD HEALTH ORGANIZATION (WHO).

Comments and contributions for publication in the WER Sri Lanka are welcome. However, the editor reserves the right to accept or reject items for publication. All correspondence should be mailed to The Editor, WER Sri Lanka, Epidemiological Unit, P.O. Box 1567, Colombo or sent by E-mail to chepid@sltnet.lk. Prior approval should be obtained from the Epidemiology Unit before publishing data in this publication

ON STATE SERVICE

Dr. P. PALIHAWADANA
 CHIEF EPIDEMIOLOGIST
 EPIDEMIOLOGY UNIT
 231, DE SARAM PLACE
 COLOMBO 10